

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS DTIC ELECTE		
2a. SECURITY CLASSIFICATION AUTHORITY SEP 11 1989			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for Public Release. Distribution Unlimited.		
2b. DECLASSIFICATION/CONTINUATION ED DATE MB B			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6b. OFFICE SYMBOL (If applicable)			7a. NAME OF MONITORING ORGANIZATION		
Texas A&M University			Texas A&M Research Foundation		
6c. ADDRESS (City, State, and ZIP Code) Department of Chemistry College Station, Texas 77843			7b. ADDRESS (City, State, and ZIP Code) Box 3578 College Station, Texas 77843		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (If applicable) ONR	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-89-J-1214		
8c. ADDRESS (City, State, and ZIP Code) 800 North Quincy Street, Code 1513:CF Arlington, VA 22217-5000			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.
			WORK UNIT ACCESSION NO.		
11. TITLE (Include Security Classification) Macrocycles Containing Tin. Ditopic, Tricyclic, Lewis Acidic Hosts with Four Binding Sites					
12. PERSONAL AUTHOR(S) Michael T. Blanda, Martin Newcomb*					
13a. TYPE OF REPORT Technical		13b. TIME COVERED FROM TO		14. DATE OF REPORT (Year, Month, Day) 1989, August 30	
				15. PAGE COUNT 8	
16. SUPPLEMENTARY NOTATION Published in <u>Tetrahedron Lett.</u>					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	macrocycle macrotricyclic Lewis acid chloride		
			complexation NMR		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) A new class of Lewis acidic hosts is described. The hosts are macrotricyclic species containing four tin atoms that serve as the bridgeheads of the macrotricyclic and contain an electron withdrawing group to impart Lewis acidic character. Variable length polymethylene chains link the tin atoms. The preparation of two macrotricyclic host precursors and one macrotricyclic host, 1,7,13,19-tetrachloro-1,7,13,19-tetrastannatrimethylene (17.5.5.5 ^{1,13}) tetratriacene (7), are described. ¹¹⁹ Sn NMR studies of host 7 binding chloride ion in CDCl ₃ at 20 °C are reported. Fast exchange between bulk and bound chloride was observed. Host 7 binds one chloride ion strongly and shows no evidence for binding additional chloride ions. These results suggest that chloride is bound within the central cavity of the host with the multiple Lewis acidic sites acting in a cooperative manner.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL			22b. TELEPHONE (Include Area Code)		22c. OFFICE SYMBOL

OFFICE OF NAVAL RESEARCH

Grant N00014-89-J-1214

R&T Code 4135009

Technical Report No. 2

Macrocycles Containing Tin.
Ditopic, Tricyclic. Lewis Acidic Hosts with Four Binding Sites

by

Michael T. Blanda and Martin Newcomb*

Department of Chemistry
Texas A&M University
College Station, TX 77843

August 30, 1989

Prepared for publication in

Tetrahedron Letters

Reproduction in whole or in part is permitted for any purpose of
the United States Government

This document has been approved for public release
and sale; its distribution is unlimited.

Macrocycles Containing Tin.


Ditopic, Tricyclic, Lewis Acidic Hosts with Four Binding Sites

Michael T. Blanda and Martin Newcomb*


Department of Chemistry, Texas A&M University, College Station, Texas, 77843, USA

Summary: Preparation of the title class of anion-binding hosts is described, and the chloride binding properties of one host are reported.

The complexation of anions in organic media by multidentate, neutral, Lewis acidic hosts is a relatively young field of study.¹ Our contributions in this area have involved the development of macrocyclic² (1) and macrobicyclic³ (2) hosts that contain Lewis acidic tin atoms as the binding sites. Low size selectivity was observed in chloride binding by macrocyclic hosts containing two binding sites,^{2b} but substantial size selectivity in binding fluoride, chloride, and bromide anions was found in two-site macrobicycles.^{3b,d} Solution ¹¹⁹Sn NMR studies of bicyclic hosts binding anions in halogenated solvents^{3b,d} as well as X-ray crystallographic and solid state ¹¹⁹Sn NMR studies of bicycle-anion complexes^{3c} demonstrate that anions are held in the cavities of the hosts; the selectivity apparently originates from the fit of the encrypted anion in the cavity. It was expected that the incorporation of additional binding sites into relatively rigid host superstructures would result in both stronger binding and higher selectivity. In this letter, we report examples of macrotricyclic species containing four tin atoms and studies of chloride complexation by one ditopic host that exhibits the expected enhancement in anion binding strength.




1: n = 8, 10, 12



2: X=Cl or Br
n=6,7,8,10,12


Our approach to the synthesis of tin-containing hosts involves sequential addition of the linking polymethylene chains. Thus, one reactive site per tin atom was created by selective HX cleavage of phenyl groups from a bis(triphenylstannyl)alkane, and the resulting bis(halodiphenylstannyl)alkane was converted to a macrocycle by reaction with a di-Grignard reagent under relatively high dilution condi-

tions.^{2a} Repeating the sequence on macrocycles containing linking polymethylene chains of 6, 7, 8, 10, or 12 units gave macrobicyclic compounds.^{3d} For the small macrocyclic compounds 3 and 4, reaction with a di-Grignard reagent in somewhat more concentrated solutions resulted in the formation of macrotricycles 5 and 6 as the major products isolated by preparative reverse-phase chromatography.⁴ Treatment of the tetra-tin tricycle 6 with HCl gave host 7 in essentially quantitative yield.⁵



3: $n = 4$

4: $n = 5$



5: $n=4$, $X=Ph$

6: $n=5$, $X=Ph$

7: $n=5$, $X=Cl$

The chloride binding properties of host 7 in solution were studied by ^{119}Sn NMR spectroscopy. In CDCl_3 solutions, the signal from triorganotin chlorides is observed at about δ 150, and the signal from a stannate $(\text{R}_3\text{SnCl}_2)^-$ in CDCl_3 is observed in the vicinity of δ -60.^{3d} For the macrobicyclic hosts 2 that bind only one halide ion, the signal for the complexed tin atoms was found midway between these values.^{3d} When aliquots of tetrahexylammonium chloride solution were added to a CDCl_3 solution containing host 7 and NMR spectra were recorded, the limiting chemical shift for the complex was at δ 87. The figure shows the saturation curves obtained for Bu_3SnCl , macrobicyclic 2 ($n = 12$), and host 7. The break in the saturation curve for 7 at a guest to host ratio of about 1:1 not only shows that the host forms a 1:1 complex with chloride but also indicates that the binding constant is relatively large. The asymptotic value in the saturation study of δ 87, about one fourth of the total change in chemical shift expected for conversion of stannanes to stannates, confirms that a 1:1 complex is formed. Further, the lack of change in the chemical shift in NMR spectra of 7 in the presence of excess chloride shows that any interaction between the 1:1 complex and a second chloride anion is very weak.

A single ^{119}Sn NMR signal was observed from all solutions containing host 7 and chloride which shows that the four tin atoms in the host bind the anion either simultaneously or in an NMR-fast, time averaged manner. Exchange of free host and the 1:1 complex also was quite rapid. In the absence of added chloride, the ^{119}Sn NMR spectrum of 7 was complex due to slow conformational changes in solution, and it was difficult to estimate the natural T_2 for this species. Nevertheless, using a conservative estimate of 0.0005 s for T_2 of free host 7, line shape simulations⁶ of spectra containing one and two equivalents of chloride revealed that noticeable line broadening due to exchange would have been apparent if the rate constant for exchange was $1 \times 10^6 \text{ s}^{-1}$ or less at 20 °C.

Availability Codes

Dist

Avail and/or
Special

A-1




Figure. Titration Curves for Bu_3SnCl (□), bicycle 2 ($n = 12$) (○), and host 7 (Δ) in CDCl_3 .

The chloride binding behavior of macrotricyclic host 7 is unique in comparison to that seen with the macrocyclic and macrobicyclic species we have previously studied. Specifically, fast exchange of free host and complex was observed with the macrocyclic hosts 1, but these species showed only limited cooperative effects of the binding sites and bound two chloride anions strongly.^{2b} On the other hand, for the bicyclic hosts 2 that formed 1:1 complexes with chloride, exchange of free host and complex by dissociation of the bound chloride from the cavity tended to be a slow process;^{3d} indeed, in the bicycle that bound chloride most strongly, the rate constant for dissociation of chloride from the cavity of the complex was on the order of $1 \times 10^4 \text{ s}^{-1}$, and the chloride selectivity for hosts 2 was found to originate predominantly in the rates of dissociation of the complexes.^{3d} That the tricyclic host 7 forms a 1:1 complex with chloride while rapid equilibration between free host and the complex is maintained suggests that the propensity for forming the 1:1 complex may result more from the cooperative binding of the guest by the multiple binding sites than from a simple structure-controlled exclusion of additional guests.

The formation constant for the 1:1 complex of 7 and chloride was determined by an iterative Hildebrand-Benesi treatment.⁷ At 20 °C, the binding constant in CDCl_3 was 500 M^{-1} . This represents an increase in binding energy in CDCl_3 solution of 2.0 kcal/mol over that found for the strongest binding bicyclic host (2, $n = 8$).^{3d} It is noteworthy that this binding constant is also substantially greater than that for Bu_3SnCl binding chloride. Previously, we observed that, although the bicyclic hosts 2 could bind chloride selectively, the largest binding constant was no greater than that of Bu_3SnCl ,^{3d} and we ascribed this behavior in part to the fact that hosts 2 apparently bind chloride with only one Lewis acidic site. The substantial increase in the binding constant for host 7 complexing chloride is additional evidence that the multiple Lewis acid sites in 7 act in a cooperative manner.

In summary, the strong binding of chloride by host 7 coupled with its propensity to form only a 1:1 complex with chloride suggest that the Lewis acidic sites in the ditopic host are properly oriented for binding an included guest. In our continuing efforts with four site, Lewis acidic hosts, we plan to vary

the length of the spacers between the two binding regions defined by the macrocyclic subunits to give a series of ditopic hosts. Given the unique chloride binding features of host 7, we expect that the macrotricycles reported here are the progenitors of a series of interesting anion and donor binding agents.

Acknowledgment: We thank the Office of Naval Research for financial support. We thank Mr. W. M. Cates for assistance in the preparation of 5 and Dr. J. H. Horner for helpful discussions.

References and Notes

- Recent examples include the following. Jung, M. E.; Xia, H. *Tetrahedron Lett.* **1988**, 29, 297-300. Wuest, J. D.; Zacharie, B. *J. Am. Chem. Soc.* **1987**, 109, 4714-4715. Katz, H. E. *Organometallics* **1987**, 6, 1134-1136. Swami, K.; Hutchinson, J. P.; Kuivila, H. G.; Zubietta, J. A. *Ibid.* **1984**, 3, 1687-1694. Gielen, M.; Jurkschat, K.; Mahieu, B.; Apers, D. *J. Organometal. Chem.* **1985**, 286, 145-151.
- (a) Azuma, Y.; Newcomb, M. *Organometallics* **1984**, 3, 9-14. (b) Newcomb, M.; Madonik, A. M.; Blanda, M. T.; Judice, J. K. *Ibid.* **1987**, 6, 145-150.
- (a) Newcomb, M.; Horner, J. H.; Blanda, M. T.; *J. Am. Chem. Soc.* **1987**, 109, 7878-7879. (b) Newcomb, M.; Blanda, M. T. *Tetrahedron Lett.* **1988**, 29, 4261-4264. (c) Newcomb, M.; Horner, J. H.; Blanda, M. T.; Squattrito, P. J. *J. Am. Chem. Soc.* **1989**, 111, 0000. (d) Blanda, M. T.; Horner, J. H.; Newcomb, M., submitted for publication.
- In a representative procedure, a solution of 4^{3d} (4 mmol) in 500 mL of THF under nitrogen was stirred at room temperature as a solution of the di-Grignard reagent from 1,5-dibromopentane in THF (500 mL, 0.012 M) was added over a period of 2 h. The mixture was stirred for an additional 6 h at room temperature. Approximately half of the THF was distilled from the reaction mixture at reduced pressure, and the remaining solution was diluted with 200 mL of ether. The mixture was washed with sat'd NH₄Cl soln, sat'd KF soln, and sat'd NaCl soln. The organic phase was dried (MgSO₄) and distilled at reduced pressure to give a residue that was purified by reverse-phase chromatography (LiChroprep® RP-18, THF/MeOH elution) to give 6. Recrystallization from THF/MeOH gave 6 (mp > 220 °C) in 10% yield. Product 5 had mp 176-178 °C.
The macrotricycles were characterized as follows. Compound 5: ¹³C NMR (50 MHz, CDCl₃, ref. Me₄Si) δ 9.42, 10.29, 29.43, 31.43, 128.0, 128.2, 136.4, 142.5; ¹¹⁹Sn NMR (149 MHz, CDCl₃, ref. Me₄Sn) δ -44.2; Anal. for C₄₈H₆₈Sn₄, calcd: C, 51.48; H, 6.12; found: C, 51.90; H, 6.14. Compound 6: ¹³C NMR (50 MHz, CDCl₃, ref. Me₄Si) δ 10.84, 11.68, 27.0, 27.1, 37.3, 40.9, 128.5, 128.6, 136.7, 142.3; ¹¹⁹Sn NMR (149 MHz, CDCl₃, ref. Me₄Sn) δ -44.2; Anal. for C₅₄H₈₀Sn₄, calcd: C, 53.89; H, 6.70; found: C, 53.73; H, 6.81. (Analyses performed by Galbraith Laboratories, Inc.)
The structure of 5 was confirmed by X-ray crystallography; the results (submitted for the referees) will be published. Crystals of 6 were not suitable for X-ray analysis. The structure of 6 was deduced from (1) the analogy in the reactions forming 5 and 6, (2) the analogy in HPLC retention times of 5 and 6, (3) the high symmetry of 6 indicated by NMR spectroscopy (one ¹¹⁹Sn signal, ten ¹³C signals), (4) analysis, and (5) conversion of 6 to 7 which displayed similar high symmetry by NMR spectroscopy.
- Tetraphenyl tricycle 6 was converted to host 7 by treatment with a standardized solution of HCl in CH₂Cl₂ by the same method as reported for the cleavage reactions leading to bicyclic hosts 2.^{3d} The crude product was recrystallized from CHCl₃/hexanes to give 7 (mp > 220 °C) in 98% yield. The product had the following spectral properties: ¹³C NMR (50 MHz, CDCl₃, ref. Me₄Si) δ 18.5, 19.0, 24.7, 25.8, 37.0, 36.9; ¹¹⁹Sn NMR (149 MHz, CDCl₃, ref. Me₄Sn) δ 158.3 (major), 162.5 (minor). The ¹¹⁹Sn NMR spectra in the presence of chloride contained one sharp peak, see text. Crystals of 7 were not suitable for X-ray analysis.
- ¹¹⁹Sn NMR spectra were obtained at 149.5 MHz. The spectra were simulated with a two site exchange model; see Sandström, J. *Dynamic NMR Spectroscopy*, Academic: London, 1982; pp 12-15.
- The NMR adaptation of the Hildebrand-Benesi treatment⁸ and the iterative program have been described.^{3d} ¹¹⁹Sn NMR spectra of five solutions of host 7 and tetrahexylammonium chloride in CDCl₃ were obtained and analyzed. The resulting calculated chemical shift for the 1:1 complex was δ 87, the same as that found experimentally in the saturation study.
- Connors, K. A. *Binding Constants, The Measurement of Molecular Complex Stability*; Wiley-Interscience: New York, 1987; pp 24-28.